

CONTENTS

EDITOR'S DESK

84A Reversing wetlands history

Mark Anderson-Wilk, editor

YOUR LETTERS

86A Readers' forum

HOME FRONT

88A The promise and peril of biofuels

Craig A. Cox, executive director of the Soil and Water Conservation Society

VIEWPOINTS

90A Partnering to achieve conservation goals

Edward B. Knipling, administrator of the USDA Agricultural Research Service

93A Making every acre count for conservation

Arlen L. Lancaster, chief of the USDA Natural Resources Conservation Service

CONSERVATIONIST PROFILE

96A Jonny Harris, Greenview Farms, Georgia

CONSERVATION IN PRACTICE

97A Attitudes of Great Plains producers about best management practices, conservation programs, and water quality

Craig M. Smith, Jeffrey M. Peterson, and John C. Leatherman

CONSERVATION EDUCATION

104A Streamside science: Tailoring watershed education to meet the needs of teachers

Nancy O. Mesner and Andréé D. Walker

FEATURE

110A A decade of advances in cover crops: Cover crops with limited irrigation can increase yields, crop quality, and nutrient and water use efficiencies while protecting the environment

Jorge A. Delgado, Merlin A. Dillon, Richard T. Sparks, and Samuel Y.C. Essah



110A

Cover crops with limited irrigation can increase yields, crop quality, and nutrient and water use efficiencies while protecting the environment.



On the Cover

Satellite image of Napa Valley courtesy of DigitalGlobe. NCDC Imaging provides GIS and remote sensing applications for natural resource management using DigitalGlobe images.



TECH TRANSFER BRIEFING

118A

Using compost to encourage vegetation and limit erosion on mountain road cuts

Matthew J. Curtis, Mark E. Grismer, and Victor P. Claassen

119A

Impact of land use fragmentation and larger herd sizes on manure recycling energy and cost

William L. Bland

119A

Improving riparian buffer performance

Mark D. Tomer, Thomas B. Moorman, John L. Kovar, David E. James, and Michael R. Burkart

120A

Predicting wind erosion and windblown dust emissions at the regional scale to guide strategic conservation targeting

Guanglong Feng and Brenton Sharratt

121A

Reducing the impacts of mechanical thinning in a southwestern forest

Douglas S. Cram, Terrell T. Baker, Alexander G. Fernald, Anthony Madrid, and Bob Rummer

IN THE NEWS

122A

Conservation news you can use

APPLIED RESEARCH

321

Scaling from field to region for wind erosion prediction using the Wind Erosion Prediction System and geographical information systems

G. Feng and B. Sharratt

329

Spatial patterns of sediment and phosphorus in a riparian buffer in western Iowa

M.D. Tomer, T.B. Moorman, J.L. Kovar, D.E. James, and M.R. Burkart

338

Effect of compost incorporation on infiltration capacity and erosion from a decomposed granite road cut

M.J. Curtis, M.E. Grismer, and V.P. Claassen

345

Theoretical analysis of manure transport distance as a function of herd size and landscape fragmentation

K.D. Bartelt and W.L. Bland

353

Are cover crops being used in the US corn belt?

J.W. Singer, S.M. Nusser, and C.J. Alf

359

Mechanical thinning impacts on runoff, infiltration, and sediment yield following fuel reduction treatments in a southwestern dry mixed conifer forest

D.S. Cram, T.T. Baker, A.G. Fernald, A. Madrid, and B. Rummer



Printed on 100% recycled paper
with vegetable-based inks.

Published by
SOIL AND WATER CONSERVATION SOCIETY
945 SW Ankeny Road
Ankeny, IA 50023 USA
phone 515-289-2331
fax 515-289-1227
www.swcs.org

EXECUTIVE DIRECTOR
Craig A. Cox

EDITOR
Mark Anderson-Wilk

EDITORIAL ASSISTANT
Jacqueline Smith

RESEARCH EDITOR
Jorge Delgado, USDA Agricultural Research Service

ASSOCIATE EDITORS
Mahdi Al-Kaisi, Iowa State University
Francisco Arriaga, USDA Agricultural Research Service
James Ascough II, USDA Agricultural Research Service
Grant Cardon, Utah State University
Adrian Chappell, University of Salford, Manchester, UK
Zhibao Dong, Chinese Academy of Sciences
Moustafa Elrashidi, USDA Natural Resources
Conservation Service
Ali Fares, University of Hawaii
Bradley King, USDA Agricultural Research Service
Peter Kleinman, USDA Agricultural Research Service
Kokoasse Kpombrekou-A, Tuskegee University
Birl Lowery, University of Wisconsin–Madison
Loretta Lynch, University of Maryland
Maurice Mausbach, USDA Natural Resources
Conservation Service
Laura M.J. McCann, University of Missouri
Guy Mehuys, McGill University
Kenneth Potter, USDA Agricultural Research Service
Mary Stromberger, Colorado State University
John White, Louisiana State University
John Williams, USDA Agricultural Research Service
Wanhong Yang, University of Guelph
Ted Zobeck, USDA Agricultural Research Service

EDITORIAL ADVISORS
Lynn Betts, USDA Natural Resources Conservation
Service
Warren Busscher, USDA Agricultural Research Service

BOARD OF DIRECTORS
Peggie James, President & West North Central Director
Don Wysocki, Vice-President & Western Director
Jean Steiner, Secretary & At Large Director
Ira Linville, Treasurer & South Central Director
Robert Abercrombie, Southwestern Director
Ashley Andrews, Student Director
Joseph Arbour, Canada Director
Theo Dillaha, Southeastern Director
Clark Gantzer, At Large Director
Jerry Hatfield, At Large Director
James Hotaling, Northeastern Director
Mary Miller, Northern Plains Director
Gary Steinhardt, East North Central Director

The *Journal of Soil and Water Conservation* (ISSN 0022-4561) is published bimonthly by the Soil and Water Conservation Society. © 2007 by the Soil and Water Conservation Society. All rights reserved.

For subscriptions or membership, please visit www.swcs.org or call 1-800-THE-SOIL (1-800-843-7645).
POSTMASTER: Send address changes to Journal of Soil and Water Conservation, 945 SW Ankeny Road, Ankeny, IA 50023. Periodicals postage paid at Ankeny, IA, and additional mailing offices.

Submit comments, contributions, advertising queries, and requests for reprints and permissions to the editor at pubs@swcs.org. Articles appearing in the research section have undergone peer review. The Soil and Water Conservation Society assumes no responsibility for statements and opinions expressed by contributors.

Readers' forum

THINK LIKE A ROOT, THINK LIKE A RIVER

The editor's review of earth scientist David R. Montgomery's book, *Dirt: The Erosion of Civilizations*, followed by Leonard C. Johnson's letter concerning circular citations of "affordable rates of soil loss" (both in the May/June 2007 issue), provoked me to express my concern about the emphases that still appear to dominate thinking about soil conservation.

For a long time I have considered that progress in conservation of soil and water, as applied to agriculture at least, is commonly held back by a "log-jam" in thinking caused by the following:

1. A predominantly geologic/physical/chemical emphasis in describing soil. Although Hans Jenny listed organisms among the soil-forming factors, their significant contributions appear very often to have been given minimal attention.
2. A preoccupation with net rate of erosional soil loss on the assumption that severity of loss of soil productivity is primarily and in some way directly caused by this. It needs to be asked *why* runoff and erosion occur and whether hindering downslope soil movement by physical and/or vegetative methods adequately addresses the joint problems of decline in productivity (or the rising costs of maintaining it) on the one hand and of worsening hydrologic conditions of stream flow and groundwater recharge on the other.

Agriculture's outputs depend primarily on plants' production and therefore on the functioning of their root systems. A root's top-down perception of soil as a medium in which to grow and function is significantly different in emphasis from a geologist's bottom-upward perception of soil as an upper layer of the earth's crust modified by weathering.

Roots cannot function effectively if the medium they penetrate is inimical and if it does not retain sufficient available water. River flows cannot be suave throughout the year if rainwater does not infiltrate, and the consequent runoff induced by soil

impermeabilization heightens flood peaks in rainy seasons and represents a loss of potential volume of flow in dry seasons.

The entry of water into soil, benefiting both plant production and river regimes, depends on the ongoing presence of an adequate porosity throughout a soil profile. In this regard, the interstitial spaces between soil mineral particles and aggregates may be equally—or more—significant to the dynamics of root-growth and water movement as are the solid components of the soil. The loss of this porosity, whether due to compaction, to pulverization or to collapse following loss of organic matter, represents a loss of soil's usefulness. Demolishing a building in situ results in not just a pile of all the physical rubble but the loss of the spaces—rooms, corridors, kitchens, etc.—in which interesting dynamic things formerly occurred and which gave the building its usefulness. Even the possibility of "recycling" the rubble disappears if it is carted away. So it is with soil.

Nevertheless, the porosity of formerly productive soil, once damaged in this way, is capable of being repeatedly regenerated by soil-inhabiting organisms, including plants, fungi, and animals, interacting over time with the other chemical, physical and hydric factors of soil, which together determine the characteristics of its productivity. Such regeneration can occur as much in the root zones of disturbed but well-managed soils as in their undisturbed conditions, as for instance in native forests and grasslands.

It is apparent that soil-inhabiting organisms make their own habitats from the surface downward, as indicated by the photo of a roadside cutting in Malawi. I infer that leachates from the biotic transformations of organic matter may be contributing to the release of nutrient ions from the mineral particles in the profile. If so, this suggests that useful soil is probably forming—and can be formed when fostered appropriately—much more quickly from the top downward than is assumed behind the "mini-miasma of circular and interlocking citations" criticized by Leonard C. Johnson.



Roadside cutting in Malawi.
Photo by T. Francis Shaxson.

Striking and widespread examples of positive improvements in the maintenance of soil productivity can rightfully be attributed to the spread of well-managed rotational zero-tillage systems that achieve "top-down" enhancement. In such systems the maintenance of a crop-residue cover over the soil surface not only buffers against high-energy rainfall impact but also provides a regular supply of organic substrate for biotic activity in the soil, with several beneficial consequences. These include nutrient recycling and ongoing formation and re-formation of varied soil porosity, with attendant entry, retention, and movement of water. Following elimination of tillage from these production systems, soil organic matter can accumulate (benefiting the soil biota) rather than decline, as formerly happened due to excessive aeration, more-rapid transformation of organic matter by the soil biota, and a net loss of carbon, as its dioxide, to the atmosphere.

Herein lie the biologic roots of sustainability, which has kept much land productive over centuries, despite any past and present periods of people's mismanagement of their soil resources.

It is useful to "think like a root, think like a river" to perceive what characteristics of soil are needed so that landscapes can continue producing the vegetation and water on which, ultimately, we all depend. In this scenario, loss of soil porosity is not only a primary cause of damage to the root environment but also predisposes soils to suffer runoff and erosion as a consequence of such damage.

Understanding and managing the integrated functioning of the carbon cycle with the hydrologic cycle and all the ecological consequences therefrom deserve much

greater attention now, in our attempts to maintain soils' productivity and to secure water supplies, than the circular and less-than-fruitful debates and experiments about "affordable rates of soil loss."

The heresies uttered above have many implications for priorities in research, extension, training, policy making, funding, and for the sustainability and stability of farming in a changing world.

T. Francis Shaxson

Dorset, UK

Hugh Hammond Bennett Award
recipient, 1995

THE QUEST FOR ANSWERS LEADS TO MORE QUESTIONS

We would like to share a bit of early soil erosion research history and consider its past and future implications.

For many, the first recognition of the consequences of soil misuse came during America's Great Dust Bowl of the 1930s. Hugh Hammond Bennett was well aware of soil ruination and had written steadily about it throughout the 1920s; his warnings were not heeded. In 1930, largely in response to Bennett's soil conservation campaign, Texas Representative J.P. Buchanan attached an amendment to the US appropriations bill authorizing the USDA to establish a series of soil erosion experiment stations.

The recognition of the consequences of soil erosion had come even earlier as a surprise to an undergraduate student. In 1915, Rensselaer W. McClure, working under the guidance of Professor Merritt F. Miller at the University of Missouri–Columbia, was directed to measure the rainfall and runoff in small agronomic plots. After the first rain event, McClure, wanting to correctly measure runoff, asked Miller what to do with all the mud in the barrel used to collect runoff. Neither McClure nor Miller had expected the magnitude of soil erosion that occurred on soil cultivated for annual agronomic crops. Russell M. Vifquain, a graduate student, developed four plots in 1916 to evaluate soil moisture and runoff. Because of McClure's mud and Vifquain's study, in 1917 Frank L. Duley under the

direction of Miller installed the first seven soil erosion plots in Columbia. Bennett used the results from the Missouri erosion plots to lobby Congress to establish erosion experiment stations around the United States. Miller said of this early work, "This is an example of a simple investigation, developed somewhat accidentally that provided much needed information at a time when important use could be made of it."

Great soil and water conservation benefits have been produced from results of these erosion experiment stations. Of primary importance was the development of the Universal Soil Loss Equation (USLE) and the many models derived from this work. The importance of their data in developing effective conservation plans that tailor conservation practices to the land to reduce erosion to tolerable levels cannot be understated. Nationally, USLE-estimated data are used to produce the National Resource Inventory (NRI) estimates of conservation needs.

As great as this work is, we worry a bit that all who use this information may not realize its limitations. Soil loss estimates from USLE models only estimate the long-term average annual sheet (interrill) and rill from a standard plot less than 100 feet long. What would the NRI maps of soil erosion look like if estimates of ephemeral and gully erosion formation and erosion estimation were also included? What would NRI map estimates of actual (real time) sediment transport to surface waters show? We would like to find out and believe other conservationists would as well.

New work is being done by researchers at the Massachusetts Institute of Technology who are developing channel-hillslope integrated landscape development models that simulate erosion and sedimentation in river basins. Ongoing work at Vanderbilt University may dispel misconceptions about how rain splash transport occurs. Imagine that—something we thought Ellison had described more than 50 years ago. The more we learn, the more we realize that we have even more to learn.

In his 2004 SWCS conference keynote address, "Soil Conservation and SWCS: A Forty-Year Retrospective," Max Schnepf identified challenges that currently confront our community. He opined that while "our

movement has enjoyed a reasonable degree of political success, success on the technical side is less apparent." He suggested that technical issues will "demand the attention of NRCS and other institutions that traditionally have possessed responsibility for conservation technology development and transfer." He also argued, "To effectively address this issue will likely require added commitments from other partners and constituencies in position to help." Perhaps SWCS should consider this challenge and work to sponsor an international effort across all disciplines, comparing the advantages of the many approaches for estimating soil and water loss including use of such models as RUSLE2, WEPP, and EUROSEM and those from geologic disciplines. Such an effort would be beneficial to advancing our societal goals to "improve ways to use land that sustains its productive capacity and protects the environment." This could strengthen the conservation connection between scientists, conservationists, and landowners.

In 2004, Jocelyn Kaiser wrote in *Science*, "Soil degradation in all its nefarious forms is not a prelude to mass starvation, as analysts once feared. Nevertheless, it is eroding crop yields and contributing to malnourishment in many corners of the globe." Conservation of soil productivity and protection of environmental quality remain our primary goals. While much has been learned from work based on results from the soil erosion experiment stations, our knowledge of the processes is not yet perfect, and thus our conservation efforts are not as effective as they might be. Recognizing this problem is a step in the right direction for improved soil and water conservation, improved productivity, and a better environment.

Clark Gantzer

Professor, Department of Soil,
Environmental and Atmospheric Sciences,
University of Missouri, and At Large
Director, SWCS Board of Directors

Steve Anderson

Professor and Chair, Department of Soil,
Environmental and Atmospheric Sciences,
University of Missouri
Columbia, Missouri